




# Enzyme Reaction Rates

Enzymes are sensitive molecules. They often have a narrow range of conditions under which they operate properly. At low temperatures there is little activity. As temperature is increased, so does the enzyme activity until the point is reached when the temperature is so high it damages the protein (**denaturation**). This causes the enzyme to stop working. Extremes in acidity can also cause the protein structure of enzymes to denature. Poisons often work by

causing enzymes to cease functioning. Cofactors such as vitamins and trace elements are required for many enzymes to function.

In the four graphs below, the *rate of reaction* or degree of *enzyme activity* is plotted against each of four factors that affect enzyme performance. Answer the questions that relate to each graph:




### 1. Enzyme Concentration

(a) Describe the change in the rate of reaction when the enzyme concentration is increased (assuming there is plenty of the substrate present):

Increases reaction rate

(b) Suggest how a cell may vary the amount of enzyme present in a cell:

By manufacturing more or less enzyme (↑ or ↓ protein synthesis)




### 2. Substrate Concentration

(a) Describe the change in the rate of reaction when the substrate concentration is **increased** (assuming a fixed amount of enzyme and ample cofactors):

Increases to a point, then levels off

(b) Explain why the rate changes the way it does:

eventually all enzymes are occupied so rate cannot ↑ more.



### 3. Temperature


Higher temperatures speed up all reactions, but few enzymes can tolerate temperatures higher than 50–60°C. The rate at which enzymes are **denatured** (change their shape and become inactive) increases with higher temperatures.

(a) Describe what is meant by an *optimum temperature* for enzyme activity:

temp. at which enzyme activity is at its maximum

(b) Explain why most enzymes perform poorly at low temperatures:

bc chem rxns happen slowly (low energy = slow particles = few collisions)



### 4. Acidity (pH)

Like all proteins, enzymes are **denatured** by *extremes* of **pH** (very acid or alkaline). Within these extremes, most enzymes are still influenced by pH. Each enzyme has a preferred pH range for optimum activity.

(a) State the optimum pH for each of the enzymes:

Pepsin: 1-2 Trypsin: 7.5-8.2 Urease: 6.5-7

(b) Pepsin acts on proteins in the stomach. Explain how its optimum pH is suited to its working environment:

Stomach is an acidic environment, so ideal for pepsin




# 2 Enzyme Cofactors and Inhibitors

Enzyme activity is often influenced by the presence of other chemicals. Some of these can enhance an enzyme's activity. Called **cofactors**, they are a nonprotein component of an

enzyme and may be organic molecules (called **coenzymes**) or inorganic ions (e.g.  $Ca^{2+}$ ,  $Zn^{2+}$ ). Enzymes may also be deactivated, temporarily or permanently, by enzyme **inhibitors**.


### Types of Enzyme

Nearly all enzymes are made of protein, although RNA has been demonstrated to have enzymatic properties. Some enzymes consist of just protein, while others require the addition of extra components to complete their catalytic properties. These may be permanently attached parts called **prosthetic groups**, or temporarily attached pieces (**coenzymes**) that detach after a reaction, and may participate with another enzyme in other reactions.

| Protein-only Enzymes  | Conjugated Protein Enzymes   |  |
|---|--|--|
|  <p>Active site<br/>enzyme</p> |  <p>Active site<br/>Apoenzyme</p>           |  <p>Active site<br/>Apoenzyme<br/>Prosthetic group is more or less permanently attached<br/>Coenzyme becomes detached after the reaction</p> |
| <p>Enzyme comprised of just protein<br/>eg. Lysozyme</p>  | <p><b>Prosthetic Group Required</b><br/>Contains apoenzyme (protein) plus a prosthetic group<br/>e.g. Flavoprotein + FAD</p> | <p><b>Coenzyme Required</b><br/>Contains apoenzyme (protein) plus a coenzyme (non-protein)<br/>e.g. Dehydrogenases + NAD</p>   |


---

### Reversible Enzyme Inhibitors




Substrate  
Enzyme  
Good fit

**No inhibition**




Competitive inhibitor blocks the active site

**Competitive inhibitor**



The substrate binds to the active site but slows the speed of reaction


**Noncompetitive inhibitor**



The substrate cannot bind to the active site  
Active site is distorted

**Allosteric Enzyme inhibitor**

### Irreversible Inhibitors (Poisons)



Substrate  
Active site  
Mercury acts as an inhibitor and blocks the active site  
Papain enzyme

Certain **heavy metals** bind tightly and permanently to the active sites of enzymes, destroying their catalytic properties. Examples of toxic heavy metals include: *mercury* (Hg), *cadmium* (Cd), *lead* (Pb), and *arsenic* (As). They are generally non-competitive inhibitors, although an exception is mercury that deactivates the enzyme papain. They are retained in the body, and lost slowly.

- Describe the general role of **cofactors** in enzyme activity: non-protein

---

- Heavy metals** can be very toxic to life forms.
  - Name 4 heavy metals that are toxic to humans: \_\_\_\_\_
  - Explain in general terms why these heavy metals are toxic: \_\_\_\_\_

---

- There are many enzyme inhibitors that are not heavy metals (eg. those found in some pesticides)
  - Name a **common poison** that is an enzyme inhibitor, but *not a heavy metal*.  
\_\_\_\_\_
  - Try to find out how this poison works in interfering with enzyme function. Briefly describe which enzyme it affects:  
\_\_\_\_\_

---

- Explain the difference between **competitive** and **noncompetitive** inhibition: \_\_\_\_\_

---

- Explain how **allosteric inhibitors** differ from other noncompetitive inhibitors: \_\_\_\_\_

---